

# Modeling and Verification of Probabilistic Systems

Joost-Pieter Katoen

Lehrstuhl für Informatik 2  
Software Modeling and Verification Group

<http://moves.rwth-aachen.de/teaching/ws-1819/movep18/>

November 5, 2018

# Overview

- 1 Introduction
- 2 Qualitative PCTL
- 3 Computation Tree Logic
- 4 CTL versus qualitative PCTL
- 5 Fair CTL versus qualitative PCTL
- 6 Repeated reachability and persistence
- 7 Summary

# Summary of previous lecture

interval  
path  
formula

## Probabilistic CTL

- ▶ Allows for path properties, such as (bounded) until and next.
- ▶ State formulas include propositional logic + the operator  $\mathbb{P}_J(\varphi)$
- ▶  $s \models \mathbb{P}_J(\varphi)$  if the probability of all paths starting in  $s$  fulfilling  $\varphi$  is in  $J$
- ▶ Model checking is done by a recursive descent over the formula
- ▶ This yields a polynomial-time algorithm (linear in  $|\Phi|$ ).

# Summary of previous lecture

## Probabilistic CTL

- ▶ Allows for path properties, such as (bounded) until and next.
- ▶ State formulas include propositional logic + the operator  $\mathbb{P}_J(\varphi)$
- ▶  $s \models \mathbb{P}_J(\varphi)$  if the probability of all paths starting in  $s$  fulfilling  $\varphi$  is in  $J$
- ▶ Model checking is done by a recursive descent over the formula
- ▶ This yields a polynomial-time algorithm (linear in  $|\Phi|$ ).

# Aim of this lecture

or  $> 0$

- ▶ Is PCTL, restricted to  $\mathbb{P}_{=1}(\varphi)$ , equally expressive as CTL?
- ▶ What is the expressive power of PCTL? Can repeated reachability be expressed?

# Aim of this lecture

- ▶ Is PCTL, restricted to  $\mathbb{P}_{=1}(\varphi)$ , equally expressive as CTL?
- ▶ What is the expressive power of PCTL? Can repeated reachability be expressed?

## Set up of this lecture

1. Qualitative PCTL versus CTL.
2. Qualitative PCTL versus CTL with fairness.
3. Repeated reachability probabilities in PCTL.


# Overview

- 1 Introduction
- 2 Qualitative PCTL**
- 3 Computation Tree Logic
- 4 CTL versus qualitative PCTL
- 5 Fair CTL versus qualitative PCTL
- 6 Repeated reachability and persistence
- 7 Summary

# PCTL syntax

## Probabilistic Computation Tree Logic: Syntax

PCTL consists of state- and path-formulas.





# PCTL syntax

## Probabilistic Computation Tree Logic: Syntax

PCTL consists of state- and path-formulas.

- ▶ PCTL *state formulas* over the set  $AP$  obey the grammar:

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \mathbb{P}_J(\varphi)$$

where  $a \in AP$ ,  $\varphi$  is a path formula and  $J \subseteq [0, 1]$  is an interval.

- ▶ PCTL *path formulae* are formed according to the following grammar:

$$\varphi ::= \bigcirc \Phi \mid \Phi_1 \mathsf{U} \Phi_2 \mid \Phi_1 \mathsf{U}^{\leq n} \Phi_2$$

where  $\Phi$ ,  $\Phi_1$ , and  $\Phi_2$  are state formulae and  $n \in \mathbb{N}$ .

# Qualitative PCTL

## Qualitative PCTL

State formulae in the *qualitative fragment* of PCTL (over  $AP$ ):

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \mathbb{P}_{>0}(\varphi) \mid \mathbb{P}_{=1}(\varphi)$$

where  $a \in AP$ , and  $\varphi$  is a path formula

# Qualitative PCTL

## Qualitative PCTL

State formulae in the *qualitative fragment* of PCTL (over  $AP$ ):

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \mathbb{P}_{>0}(\varphi) \mid \mathbb{P}_{=1}(\varphi)$$

where  $a \in AP$ , and  $\varphi$  is a path formula formed according to the grammar:

$$\varphi ::= \bigcirc \Phi \mid \Phi_1 \cup \Phi_2.$$

### Remark

The probability bounds  $= 0$  and  $< 1$  can be derived:

$$\mathbb{P}_{=0}(\varphi) \equiv \neg \mathbb{P}_{>0}(\varphi) \quad \text{and} \quad \mathbb{P}_{<1}(\varphi) \equiv \neg \mathbb{P}_{=1}(\varphi)$$

So, in qualitative PCTL, there is no bounded until, and only  $> 0$ ,  $= 0$ ,  $> 1$  and  $= 1$  are allowed thresholds.

# Qualitative PCTL

## Qualitative PCTL

State formulae in the *qualitative fragment* of PCTL (over  $AP$ ):

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \mathbb{P}_{>0}(\varphi) \mid \mathbb{P}_{=1}(\varphi)$$

where  $a \in AP$ , and  $\varphi$  is a path formula formed according to the grammar:

$$\varphi ::= \bigcirc \Phi \mid \Phi_1 \cup \Phi_2.$$

$\Pr > 0$

## Examples

$$\mathbb{P}_{=1}(\underbrace{\Diamond \mathbb{P}_{>0}(\bigcirc a)})$$

almost surely you eventually reach a state that has an  $a$ -successor with

# Qualitative PCTL

## Qualitative PCTL

State formulae in the *qualitative fragment* of PCTL (over  $AP$ ):

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \mathbb{P}_{>0}(\varphi) \mid \mathbb{P}_{=1}(\varphi)$$

where  $a \in AP$ , and  $\varphi$  is a path formula formed according to the grammar:

$$\varphi ::= \bigcirc \Phi \mid \Phi_1 \cup \Phi_2.$$

## Examples

$\mathbb{P}_{=1}(\Diamond \mathbb{P}_{>0}(\bigcirc a))$  and  $\mathbb{P}_{<1}(\mathbb{P}_{>0}(\Diamond a) \cup b)$  are qualitative PCTL formulas.

# Overview

- 1 Introduction
- 2 Qualitative PCTL
- 3 Computation Tree Logic**
- 4 CTL versus qualitative PCTL
- 5 Fair CTL versus qualitative PCTL
- 6 Repeated reachability and persistence
- 7 Summary

# Computation Tree Logic

[Clarke &amp; Emerson, 1981]

## Computation Tree Logic: Syntax

CTL consists of state- and path-formulas.

- CTL *state formulas* over the set  $AP$  obey the grammar:

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \exists \varphi \mid \forall \varphi$$

where  $a \in AP$  and  $\varphi$  is a path formula

$s \models \forall \varphi$  iff all paths starting in  $s$  satisfy  $\varphi$ .

# Computation Tree Logic

[Clarke &amp; Emerson, 1981]

## Computation Tree Logic: Syntax

CTL consists of state- and path-formulas.

- ▶ CTL *state formulas* over the set  $AP$  obey the grammar:

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg \Phi \mid \exists \varphi \mid \forall \varphi$$

where  $a \in AP$  and  $\varphi$  is a path formula formed by the grammar:

$$\varphi ::= \bigcirc \Phi \mid \Phi_1 \cup \Phi_2$$

*Handwritten note:*  $P_{=1}(\Diamond P_{>0}(\bigcirc a))$

### Remark

No bounded until, and only universal and existential path quantifiers.

### Examples

$\forall \Diamond \bigcirc a$  and  $\exists (\Diamond \forall a) \cup b$  are CTL formulas.



# Computation Tree Logic

[Clarke &amp; Emerson, 1981]

## Computation Tree Logic: Syntax

CTL consists of state- and path-formulas.

- ▶ CTL *state formulas* over the set  $AP$  obey the grammar:

$$\Phi ::= \text{true} \mid a \mid \Phi_1 \wedge \Phi_2 \mid \neg\Phi \mid \exists\varphi \mid \forall\varphi$$





where  $a \in AP$  and  $\varphi$  is a path formula  $\varphi ::= \bigcirc\Phi \mid \Phi_1 \cup \Phi_2$

## Intuition

- ▶  $s \models \forall\varphi$  if all paths starting in  $s$  fulfill  $\varphi$
- ▶  $s \models \exists\varphi$  if some path starting in  $s$  fulfill  $\varphi$


Question: are CTL and qualitative PCTL equally expressive? **No.**

# CTL semantics




# CTL semantics


$$\neg \forall \Diamond \phi \equiv \exists \Box \neg \phi$$




$$\neg \exists \Diamond \phi$$



$$\exists \Box \psi$$



$$\forall \Box \psi$$



# CTL semantics (1)

## Notation

$\mathcal{D}, s \models \Phi$  if and only if state-formula  $\Phi$  holds in state  $s$  of (possibly infinite) DTMC  $\mathcal{D}$ . As  $\mathcal{D}$  is known from the context we simply write  $s \models \Phi$ .

## Satisfaction relation for state formulas

The satisfaction relation  $\models$  is defined for CTL state formulas by:

$s \models a$	iff $a \in L(s)$
$s \models \neg \Phi$	iff not $(s \models \Phi)$
$s \models \Phi \wedge \Psi$	iff $(s \models \Phi)$ and $(s \models \Psi)$
$s \models \exists \varphi$	iff <b>there exists</b> $\pi \in Paths(s). \pi \models \varphi$
$s \models \forall \varphi$	iff <b>for all</b> $\pi \in Paths(s). \pi \models \varphi$

where the semantics of CTL path-formulas is the same as for PCTL

# Overview

- 1 Introduction
- 2 Qualitative PCTL
- 3 Computation Tree Logic
- 4 CTL versus qualitative PCTL**
- 5 Fair CTL versus qualitative PCTL
- 6 Repeated reachability and persistence
- 7 Summary

# CTL versus qualitative PCTL

## Equivalence of PCTL and CTL Formulae

The PCTL formula  $\Phi$  is *equivalent* to the CTL formula  $\Psi$ , denoted  $\Phi \equiv \Psi$ , if  $Sat(\Phi) = Sat(\Psi)$  for each DTMC  $\mathcal{D}$ .

$$\underbrace{\exists O a}_{\Psi} \stackrel{?}{\equiv} \underbrace{P_{>0}(O a)}_{\Phi}$$

# CTL versus qualitative PCTL

## Equivalence of PCTL and CTL Formulae

The PCTL formula  $\Phi$  is *equivalent* to the CTL formula  $\Psi$ , denoted  $\Phi \equiv \Psi$ , if  $Sat(\Phi) = Sat(\Psi)$  for each DTMC  $\mathcal{D}$ .

## Example

The simplest such cases are path formulae involving the next-step operator:

$$\mathbb{P}_{=1}(\bigcirc a) \equiv \forall \bigcirc a$$

$$\mathbb{P}_{>0}(\bigcirc a) \equiv \exists \bigcirc a$$

And for  $\exists \Diamond$  and  $\forall \Box$  we have:

$$\mathbb{P}_{>0}(\Diamond a) \equiv \exists \Diamond a$$

$$\mathbb{P}_{=1}(\Box a) \equiv \forall \Box a.$$

# CTL versus qualitative PCTL

$$(1) \mathbb{P}_{>0}(\Diamond a) \equiv \exists \Diamond a \quad \text{and} \quad (2) \mathbb{P}_{=1}(\Box a) \equiv \forall \Box a.$$

## Proof:

(1) Consider the first statement.

$\Rightarrow$  Assume  $s \models \mathbb{P}_{>0}(\Diamond a)$ .

$$\rightarrow \Rightarrow \quad \underbrace{\text{Sat}(\mathbb{P}_{>0}(\Diamond a))}_{\subseteq} \subseteq \text{Sat}(\exists \Diamond a)$$

$\Leftarrow$

$\supseteq$



# CTL versus qualitative PCTL

$$(1) \mathbb{P}_{>0}(\Diamond a) \equiv \exists \Diamond a \quad \text{and} \quad (2) \mathbb{P}_{=1}(\Box a) \equiv \forall \Box a.$$

## Proof:

(1) Consider the first statement.

$\Rightarrow$  Assume  $s \models \mathbb{P}_{>0}(\Diamond a)$ . By the PCTL semantics,  $Pr(s \models \Diamond a) > 0$ .  
Thus,  $\{\pi \in Paths(s) \mid \pi \models \Diamond a\} \neq \emptyset$ , and hence,  $s \models \exists \Diamond a$ .

$$s \in \text{Sat}(\exists \Diamond a)$$

# CTL versus qualitative PCTL

$$(1) \mathbb{P}_{>0}(\Diamond a) \equiv \exists \Diamond a \quad \text{and} \quad (2) \mathbb{P}_{=1}(\Box a) \equiv \forall \Box a.$$

## Proof:

(1) Consider the first statement.

$\Rightarrow$  Assume  $s \models \mathbb{P}_{>0}(\Diamond a)$ . By the PCTL semantics,  $Pr(s \models \Diamond a) > 0$ .  
Thus,  $\{\pi \in Paths(s) \mid \pi \models \Diamond a\} \neq \emptyset$ , and hence,  $s \models \exists \Diamond a$ .

$\Leftarrow$  Assume  $s \models \exists \Diamond a$ , i.e., there is a finite path  $\hat{\pi} = s_0 s_1 \dots s_n$  with  
 $s_0 = s$  and  $s_n \models a$ .

$\underbrace{s_0}_{=s} \quad \underbrace{s_n}_a$

# CTL versus qualitative PCTL


$$(1) \mathbb{P}_{>0}(\Diamond a) \equiv \exists \Diamond a \quad \text{and} \quad (2) \mathbb{P}_{=1}(\Box a) \equiv \forall \Box a.$$

## Proof:

(1) Consider the first statement.

$\Rightarrow$  Assume  $s \models \mathbb{P}_{>0}(\Diamond a)$ . By the PCTL semantics,  $Pr(s \models \Diamond a) > 0$ .  
Thus,  $\{\pi \in Paths(s) \mid \pi \models \Diamond a\} \neq \emptyset$ , and hence,  $s \models \exists \Diamond a$ .

$\Leftarrow$  Assume  $s \models \exists \Diamond a$ , i.e., there is a finite path  $\hat{\pi} = s_0 s_1 \dots s_n$  with  $s_0 = s$  and  $s_n \models a$ . It follows that all paths in the cylinder set  $Cyl(\hat{\pi})$  fulfill  $\Diamond a$ .



# CTL versus qualitative PCTL

$$(1) \mathbb{P}_{>0}(\Diamond a) \equiv \exists \Diamond a \quad \text{and} \quad (2) \mathbb{P}_{=1}(\Box a) \equiv \forall \Box a.$$

## Proof:

(1) Consider the first statement.

$\Rightarrow$  Assume  $s \models \mathbb{P}_{>0}(\Diamond a)$ . By the PCTL semantics,  $Pr(s \models \Diamond a) > 0$ .  
Thus,  $\{\pi \in Paths(s) \mid \pi \models \Diamond a\} \neq \emptyset$ , and hence,  $s \models \exists \Diamond a$ .

$\Leftarrow$  Assume  $s \models \exists \Diamond a$ , i.e., there is a finite path  $\hat{\pi} = s_0 s_1 \dots s_n$  with  $s_0 = s$  and  $s_n \models a$ . It follows that all paths in the cylinder set  $Cyl(\hat{\pi})$  fulfill  $\Diamond a$ . Thus:

$$Pr(s \models \Diamond a) \geq Pr_s(\underbrace{Cyl(s_0 s_1 \dots s_n)}_{=\hat{\pi}}) = \mathbf{P}(\underbrace{s_0 s_1 \dots s_n}_{=\hat{\pi}}) > 0.$$

# CTL versus qualitative PCTL

$$(1) \mathbb{P}_{>0}(\Diamond a) \equiv \exists \Diamond a \quad \text{and} \quad (2) \mathbb{P}_{=1}(\Box a) \equiv \forall \Box a.$$

## Proof:

(1) Consider the first statement.

$\Rightarrow$  Assume  $s \models \mathbb{P}_{>0}(\Diamond a)$ . By the PCTL semantics,  $Pr(s \models \Diamond a) > 0$ . Thus,  $\{\pi \in Paths(s) \mid \pi \models \Diamond a\} \neq \emptyset$ , and hence,  $s \models \exists \Diamond a$ .

$\Leftarrow$  Assume  $s \models \exists \Diamond a$ , i.e., there is a finite path  $\hat{\pi} = s_0 s_1 \dots s_n$  with  $s_0 = s$  and  $s_n \models a$ . It follows that all paths in the cylinder set  $Cyl(\hat{\pi})$  fulfill  $\Diamond a$ . Thus:

$$Pr(s \models \Diamond a) \geq Pr_s(Cyl(s_0 s_1 \dots s_n)) = \mathbf{P}(s_0 s_1 \dots s_n) > 0.$$

So, by the PCTL semantics we have:  $s \models \mathbb{P}_{>0}(\Diamond a)$ .

# CTL versus qualitative PCTL

$$(1) \mathbb{P}_{>0}(\Diamond a) \equiv \exists \Diamond a \quad \text{and} \quad (2) \mathbb{P}_{=1}(\Box a) \equiv \forall \Box a.$$

## Proof:

(1) Consider the first statement.

$\Rightarrow$  Assume  $s \models \mathbb{P}_{>0}(\Diamond a)$ . By the PCTL semantics,  $Pr(s \models \Diamond a) > 0$ .  
Thus,  $\{\pi \in Paths(s) \mid \pi \models \Diamond a\} \neq \emptyset$ , and hence,  $s \models \exists \Diamond a$ .

$\Leftarrow$  Assume  $s \models \exists \Diamond a$ , i.e., there is a finite path  $\hat{\pi} = s_0 s_1 \dots s_n$  with  $s_0 = s$  and  $s_n \models a$ . It follows that all paths in the cylinder set  $Cyl(\hat{\pi})$  fulfill  $\Diamond a$ . Thus:

$$Pr(s \models \Diamond a) \geq Pr_s(Cyl(s_0 s_1 \dots s_n)) = \mathbf{P}(s_0 s_1 \dots s_n) > 0.$$

So, by the PCTL semantics we have:  $s \models \mathbb{P}_{>0}(\Diamond a)$ .

(2) The second statement follows by duality.

# CTL versus qualitative PCTL

$$(1) \mathbb{P}_{>0}(\Diamond a) \equiv \exists \Diamond a \quad \text{and} \quad (2) \mathbb{P}_{=1}(\Box a) \equiv \forall \Box a.$$

$$(3) \mathbb{P}_{>0}(\Box a) \not\equiv \exists \Box a$$

Handwritten annotations for equation (3): A wavy line under  $\Psi$  is connected by an upward arrow to a wavy line under  $\Phi$ .


# CTL versus qualitative PCTL

$$(1) \mathbb{P}_{>0}(\Diamond a) \equiv \exists \Diamond a \quad \text{and} \quad (2) \mathbb{P}_{=1}(\Box a) \equiv \forall \Box a.$$

$$(3) \mathbb{P}_{>0}(\Box a) \not\equiv \exists \Box a \quad \text{and} \quad (4) \mathbb{P}_{=1}(\Diamond a) \not\equiv \forall \Diamond a.$$

## Example

Consider the second statement (4). Let  $s$  be a state in a (possibly infinite) DTMC. Then:  $s \models \forall \Diamond a$  implies  $s \models \mathbb{P}_{=1}(\Diamond a)$ . The reverse direction, however, does not hold. Consider the example DTMC:



$s \models \mathbb{P}_{=1}(\Diamond a)$  as the probability of path  $s^\omega$  is zero. However, the path  $s^\omega$  is possible and violates  $\Diamond a$ . Thus,  $s \not\models \forall \Diamond a$ .

Statement (3) follows by duality.




# Almost-sure-reachability not in CTL

$$P_{=1}(\bigcirc \quad P_{=1}(\bigtriangleup a)) \neq$$

$$P_{=1}(\bigtriangleup \quad P_{=1}(\bigcirc a))$$

Consider the DTMC:



$$s \models P_{=1}(\bigcirc \quad \underbrace{P_{=1}(\bigtriangleup a)})$$

$$s \not\models P_{=1}(\bigtriangleup \quad P_{=1}(\bigcirc a))$$


# Almost-sure-reachability not in CTL

## Almost-sure-reachability not in CTL

1. There is no CTL formula that is equivalent to  $\mathbb{P}_{=1}(\Diamond a)$ .
2. There is no CTL formula that is equivalent to  $\mathbb{P}_{>0}(\Box a)$ .

### Proof:

We provide the proof of 1.; 2. follows by duality:  $\mathbb{P}_{>0}(\Box a) \equiv \neg \mathbb{P}_{=1}(\Diamond \neg a)$ . By contraposition. Assume  $\Phi \equiv \mathbb{P}_{=1}(\Diamond a)$ . Consider the infinite DTMC  $\mathcal{D}_p$ :



The value of  $p$  **does** affect reachability:  $Pr(s \models \Diamond s_0) = \begin{cases} 1 & \text{if } p \leq \frac{1}{2} \\ < 1 & \text{if } p > \frac{1}{2} \end{cases}$

# Almost-sure-reachability not in CTL

There is no CTL formula that is equivalent to  $\mathbb{P}_{=1}(\Diamond a)$ .

## Proof:

$$\text{We have: } Pr(s \models \Diamond s_0) = \begin{cases} 1 & \text{if } p \leq \frac{1}{2} \\ < 1 & \text{if } p > \frac{1}{2} \end{cases}$$

Thus, in  $\mathcal{D}_{\frac{1}{4}}$  we have  $s \models \mathbb{P}_{=1}(\Diamond s_0)$  for all states  $s$ ,

# Almost-sure-reachability not in CTL

There is no CTL formula that is equivalent to  $\mathbb{P}_{=1}(\Diamond a)$ .

## Proof:

$$\text{We have: } Pr(s \models \Diamond s_0) = \begin{cases} 1 & \text{if } p \leq \frac{1}{2} \\ < 1 & \text{if } p > \frac{1}{2} \end{cases}$$

Thus, in  $\mathcal{D}_{\frac{1}{4}}$  we have  $s \models \mathbb{P}_{=1}(\Diamond s_0)$  for all states  $s$ , while in  $\mathcal{D}_{\frac{3}{4}}$ , e.g.,  $s_1 \not\models \mathbb{P}_{=1}(\Diamond s_0)$ .


# Almost-sure-reachability not in CTL

There is no CTL formula that is equivalent to  $\mathbb{P}_{=1}(\Diamond a)$ .

## Proof:

$$\text{We have: } Pr(s \models \Diamond s_0) = \begin{cases} 1 & \text{if } p \leq \frac{1}{2} \\ < 1 & \text{if } p > \frac{1}{2} \end{cases}$$

Thus, in  $\mathcal{D}_{\frac{1}{4}}$  we have  $s \models \mathbb{P}_{=1}(\Diamond s_0)$  for all states  $s$ , while in  $\mathcal{D}_{\frac{3}{4}}$ , e.g.,  $s_1 \not\models \mathbb{P}_{=1}(\Diamond s_0)$ . Hence:  $s_1 \in Sat_{\mathcal{D}_{\frac{1}{4}}}(\mathbb{P}_{=1}(\Diamond s_0))$  but  $s_1 \notin Sat_{\mathcal{D}_{\frac{3}{4}}}(\mathbb{P}_{=1}(\Diamond s_0))$ .



# Almost-sure-reachability not in CTL

There is no CTL formula that is equivalent to  $\mathbb{P}_{=1}(\Diamond a)$ .

## Proof:

$$\text{We have: } Pr(s \models \Diamond s_0) = \begin{cases} 1 & \text{if } p \leq \frac{1}{2} \\ < 1 & \text{if } p > \frac{1}{2} \end{cases}$$

Thus, in  $\mathcal{D}_{\frac{1}{4}}$  we have  $s \models \mathbb{P}_{=1}(\Diamond s_0)$  for all states  $s$ , while in  $\mathcal{D}_{\frac{3}{4}}$ , e.g.,  $s_1 \not\models \mathbb{P}_{=1}(\Diamond s_0)$ . Hence:  $s_1 \in Sat_{\mathcal{D}_{\frac{1}{4}}}(\mathbb{P}_{=1}(\Diamond s_0))$  but  $s_1 \notin Sat_{\mathcal{D}_{\frac{3}{4}}}(\mathbb{P}_{=1}(\Diamond s_0))$ . For CTL-formula  $\Phi$ —by assumption  $\Phi \equiv \mathbb{P}_{=1}(\Diamond s_0)$ —we have:

$$Sat_{\mathcal{D}_{\frac{1}{4}}}(\Phi) = Sat_{\mathcal{D}_{\frac{3}{4}}}(\Phi).$$

Hence, state  $s_1$  either fulfills the CTL formula  $\Phi$  in both DTMCs or in none of them. This, however, contradicts  $\Phi \equiv \mathbb{P}_{=1}(\Diamond s_0)$ .

## Remark



## Remark

The proof relies on the fact that the satisfaction of  $\mathbb{P}_{=1}(\Diamond a)$  for infinite DTMCs may depend on the precise value of the transition probabilities, while CTL just refers to the underlying graph of a DTMC.

## Remark

The proof relies on the fact that the satisfaction of  $\mathbb{P}_{=1}(\Diamond a)$  for infinite DTMCs may depend on the precise value of the transition probabilities, while CTL just refers to the underlying graph of a DTMC. For finite DTMCs, the previous result does **not** hold.

## Remark

The proof relies on the fact that the satisfaction of  $\mathbb{P}_{=1}(\Diamond a)$  for infinite DTMCs may depend on the precise value of the transition probabilities, while CTL just refers to the underlying graph of a DTMC. For finite DTMCs, the previous result does **not** hold.

For each finite DTMC  $\mathcal{D}$  it holds that:

$$\mathbb{P}_{=1}(\Diamond a) \equiv \underbrace{\forall((\exists \Diamond a) W a)}_{\text{true } \cup}$$

where  $W$  is the weak until operator defined by  $\Phi W \Psi = \underbrace{(\Phi \cup \Psi) \vee \Box \Phi}$ .

$$a W b \equiv a \cup b \vee \underbrace{\Box a}_{\text{weakening part}}$$

for finite

$$\text{DTMCs : } P_{=1}(\Diamond a) \equiv \forall ((\exists \Diamond a) \text{ W } a)$$

Proof  $\Pr(s \models \Diamond a) < 1 \quad \leftarrow$

$$\Leftrightarrow \Pr(s \models \Box \neg a) > 0$$

$$\Leftrightarrow \exists \hat{\pi} \in \text{Paths}^*(s). \text{ Cyl}(\hat{\pi}) \cap \Diamond a = \emptyset$$

$$\Leftrightarrow (* \text{ long-run theorem for finite DTMCs } *)$$

$$\exists \hat{\pi} \in \text{Paths}^*(s). \exists \neg a\text{-BSCC } B.$$

" $\hat{\pi}$  is a  $\neg a$ -path" and

$$s \xrightarrow{\hat{\pi}} B$$

$$\Leftrightarrow \exists \pi \in \text{Paths}(s). \left( \pi \not\models \Diamond a \wedge \right. \\ \left. \pi \not\models \Box \exists \Diamond a \right)$$

$$\Leftrightarrow s \models \exists (\neg(\text{true} \vee a) \wedge \neg \forall \Box \exists \Diamond a)$$

$$\Leftrightarrow s \models \exists (\neg((\exists \Diamond a) \vee a) \wedge \neg \forall \Box \exists \Diamond a)$$

$$\Leftrightarrow s \models \neg \forall ((\exists \Diamond a) \text{ W } a)$$

$$\Leftrightarrow s \not\models \forall ((\exists \Diamond a) \text{ W } a) \quad \square$$

## Remark

The proof relies on the fact that the satisfaction of  $\mathbb{P}_{=1}(\Diamond a)$  for infinite DTMCs may depend on the precise value of the transition probabilities, while CTL just refers to the underlying graph of a DTMC. For finite DTMCs, the previous result does **not** hold.

For each finite DTMC  $\mathcal{D}$  it holds that:

$$\mathbb{P}_{=1}(\Diamond a) \quad \equiv \quad \forall ( (\exists \Diamond a) W a )$$


where  $W$  is the weak until operator defined by  $\Phi W \Psi = (\Phi U \Psi) \vee \Box \Phi$ .

## Proof:

Exercise.

# $\forall\Diamond$ is not expressible in qualitative PCTL

1. There is no qualitative PCTL formula that is equivalent to  $\forall\Diamond a$ .
2. There is no qualitative PCTL formula that is equivalent to  $\exists\Box a$ .



# $\forall\Diamond$ is not expressible in qualitative PCTL

1. There is no qualitative PCTL formula that is equivalent to  $\forall\Diamond a$ .
2. There is no qualitative PCTL formula that is equivalent to  $\exists\Box a$ .



## Proof:

Proof of the first claim on the black board. The second claim follows by duality since  $\forall\Diamond a \equiv \neg\exists\Box\neg a$ .

Claim: there is no qualitative PCTL-formula equivalent to  $\forall \Diamond a$

---

Proof (sketch):



Claim for any qualitative PCTL-formula  $\Phi$  of nesting depth  $< n$  we have

$$\left( \begin{array}{l} s_n \models \Phi \text{ iff } t_n \models \Phi \\ \text{nd}(a) = 0 \end{array} \right.$$

$$\text{nd}(\Phi \wedge \Psi) = \max(\text{nd}(\Phi), \text{nd}(\Psi))$$

$$\text{nd}(\mathbb{P}_{\geq}(\varphi)) = 1 + \text{nd}(\varphi)$$

$$\text{nd}(0 \Psi) = \text{nd}(\Psi)$$

Proof of claim: induction on  $n$ .



By contraposition, we prove the original claim. Suppose:

$$\underbrace{\Phi}_{\substack{\text{quel.} \\ \text{PCTL}}} \equiv \forall \Diamond a$$

$$\text{let } n = \text{nd}(\Phi) + 1$$

By the claim:  $s_n \models \Phi$  iff  $t_n \models \Phi$

But  $s_n \not\models \forall \Diamond a$  whereas  $t_n \models \forall \Diamond a$

Contradiction


# Qualitative PCTL versus CTL

## Incomparable expressiveness

Qualitative PCTL and CTL have incomparable expressiveness; e.g.,  $\forall \Diamond a$  cannot be expressed in qualitative PCTL and  $\mathbb{P}_{=1}(\Diamond a)$  cannot be expressed in CTL.

# Overview

- 1 Introduction
- 2 Qualitative PCTL
- 3 Computation Tree Logic
- 4 CTL versus qualitative PCTL
- 5 Fair CTL versus qualitative PCTL
- 6 Repeated reachability and persistence
- 7 Summary



$$\forall \phi \in \neq \mathbb{P}_{=1}(\phi^a)$$

# Fairness

loops with prob. 0

## Remark

The existence of unfair computations is vital (in particular  $s_n^\omega$  in the proof of the result that  $\forall\Diamond$  is not expressible in qualitative PCTL.)

# Fairness

## Remark

The existence of unfair computations is vital (in particular  $s_n^\omega$  in the proof of the result that  $\forall\Diamond$  is not expressible in qualitative PCTL.) In fact, under appropriate **fairness** constraints, we yield  $\forall\Diamond a \equiv \mathbb{P}_{=1}(\Diamond a)$ .

## Strong fairness

Assume  $\mathcal{D}$  is a finite DTMC and that any state  $s$  in  $\mathcal{D}$  is uniquely characterized by an atomic proposition, say  $s$ .

$\bigcirc s$

$\Diamond s$

# Fairness


## Remark

The existence of unfair computations is vital (in particular  $s_n^\omega$  in the proof of the result that  $\forall\Diamond$  is not expressible in qualitative PCTL.) In fact, under appropriate **fairness** constraints, we yield  $\forall\Diamond a \equiv \mathbb{P}_{=1}(\Diamond a)$ .


## Strong fairness

Assume  $\mathcal{D}$  is a finite DTMC and that any state  $s$  in  $\mathcal{D}$  is uniquely characterized by an atomic proposition, say  $s$ . The **(strong) fairness** constraint *fair* is defined by:

$$fair = \bigwedge_{s \in S} \bigwedge_{t \in Post(s)} (\Box\Diamond s \rightarrow \Box\Diamond t).$$



Fairness  $\forall \Diamond a \neq \mathbb{P}_{=1}(\Diamond a)$



## Remark

The existence of unfair computations is vital (in particular  $s_n^\omega$  in the proof of the result that  $\forall \Diamond$  is not expressible in qualitative PCTL.) In fact, under appropriate **fairness** constraints, we yield  $\forall \Diamond a \equiv \mathbb{P}_{=1}(\Diamond a)$ .

## Strong fairness

Assume  $\mathcal{D}$  is a finite DTMC and that any state  $s$  in  $\mathcal{D}$  is uniquely characterized by an atomic proposition, say  $s$ . The **(strong) fairness** constraint *fair* is defined by:

$$fair = \bigwedge_{s \in S} \bigwedge_{t \in Post(s)} (\Box \Diamond s \rightarrow \Box \Diamond t).$$

It asserts that when a state  $s$  is visited infinitely often, then every of its direct successors is visited infinitely often too.

# Fair CTL

$\Box a$

## Fair paths

In **fair** CTL, path formulas are interpreted over **fair** infinite paths, i.e., paths  $\pi$  that satisfy

$$fair = \bigwedge_{s \in S} \bigwedge_{t \in Post(s)} (\Box \Diamond s \rightarrow \Box \Diamond t).$$

LTL formula



# Fair CTL

## Fair paths

In **fair** CTL, path formulas are interpreted over **fair** infinite paths, i.e., paths  $\pi$  that satisfy

$$fair = \bigwedge_{s \in S} \bigwedge_{t \in Post(s)} (\Box \Diamond s \rightarrow \Box \Diamond t).$$

A path  $\pi$  such that  $\pi \models fair$  is called **fair**. Let  $Paths_{fair}(s)$  be the set of fair paths starting in  $s$ .

## Fair CTL semantics

The **fair semantics** of CTL is defined by the satisfaction  $\models_{fair}$  which is defined as  $\models$  for the CTL semantics, except that:

$s \models_{fair} \exists \varphi$  iff **there exists**  $\pi \in Paths_{fair}(s). \pi \models_{fair} \varphi$

$s \models_{fair} \forall \varphi$  iff **for all**  $\pi \in Paths_{fair}(s). \pi \models_{fair} \varphi$ .

# Fairness theorem

## Qualitative PCTL versus fair CTL theorem

Let  $s$  be an arbitrary state in a finite DTMC. Then:

$$s \models \mathbb{P}_{=1}(\Diamond a) \quad \text{iff} \quad s \models_{fair} \forall \Diamond a$$

$$s \models \mathbb{P}_{>0}(\Box a) \quad \text{iff} \quad s \models_{fair} \exists \Box a$$

$$s \models \mathbb{P}_{=1}(a \cup b) \quad \text{iff} \quad s \models_{fair} \forall (a \cup b)$$

$$s \models \mathbb{P}_{>0}(a \cup b) \quad \text{iff} \quad s \models_{fair} \exists (a \cup b)$$

## Proof:

Using the fairness theorem (cf. Lecture 4): for (possibly infinite) DTMC  $\mathcal{D}$  and  $s, t$  states in  $\mathcal{D}$ :

$$Pr(s \models \Box \Diamond t) = Pr(s \models \bigwedge_{u \in Post^*(t)} \Box \Diamond u).$$

In addition, we use that from every reachable state at least one fair path starts. Similar arguments hold for infinite DTMCs (where *fair* is interpreted as infinitary conjunction.)

# Qualitative PCTL versus fair CTL

## Comparable expressiveness

Qualitative PCTL and fair CTL are equally expressive for finite Markov chains.


# Overview

- 1 Introduction
- 2 Qualitative PCTL
- 3 Computation Tree Logic
- 4 CTL versus qualitative PCTL
- 5 Fair CTL versus qualitative PCTL
- 6 Repeated reachability and persistence

## 7 Summary

$\Box \Diamond a$        $\Diamond \Box a$   
 $\text{CTL}$

$\forall \phi$        $\text{CTL}$        $\forall \phi \exists \psi$



qualitative

PCTL  $\mathbb{P}_{>0}(\psi)$

$\mathbb{P}_{=1}(\psi)$

# Almost sure repeated reachability

## Almost sure repeated reachability is PCTL-definable

For finite DTMC  $\mathcal{D}$ , state  $s \in S$  and  $G \subseteq S$ :

$$\underbrace{s \models \mathbb{P}_{=1}(\Box \mathbb{P}_{=1}(\Diamond G))}_{\text{qualitative PCTL}} \quad \text{iff} \quad \underbrace{Pr_s\{\pi \in Paths(s) \mid \pi \models \Box \Diamond G\}}_{\text{almost sure repeated reachability}} = 1.$$

# Almost sure repeated reachability

## Almost sure repeated reachability is PCTL-definable

For finite DTMC  $\mathcal{D}$ , state  $s \in S$  and  $G \subseteq S$ :

$$s \models \mathbb{P}_{=1}(\Box \mathbb{P}_{=1}(\Diamond G)) \quad \text{iff} \quad Pr_s\{\pi \in Paths(s) \mid \pi \models \Box \Diamond G\} = 1.$$

We abbreviate  $\mathbb{P}_{=1}(\Box \mathbb{P}_{=1}(\Diamond G))$  by  $\mathbb{P}_{=1}(\Box \Diamond G)$ .

### Proof:

On the blackboard.

$$s \models \mathbb{P}_{=1}(\Box \mathbb{P}_{=1}(\Diamond G)) \Rightarrow$$

$$\Pr_s \{ \pi \in \text{Paths}(s) \mid \pi \models \Box \Diamond G \} = 1$$

Proof Assume  $s \models \mathbb{P}_{=1}(\Box \mathbb{P}_{=1}(\Diamond G))$ .

As  $\mathbb{P}_{=1}(\Box a) \equiv \forall \Box a$ , we have

$$s \models \forall \Box \text{sat}(\mathbb{P}_{=1}(\Diamond G))$$

So, all states  $t \in \text{Post}^*(s)$  satisfy  $\mathbb{P}_{=1}(\Diamond G)$ .

Let BSCC  $T \subseteq \mathcal{S}$  such that  $\Pr(s \models \Diamond T) > 0$ .

Then  $s \models \Diamond t$  for each  $t \in T$  and  $t \models \mathbb{P}_{=1}(\Diamond G)$ .

As  $\text{Post}^*(t) = T$  for every  $t \in T$ , so  $T \cap G \neq \emptyset$ .

Hence, for almost all paths  $\pi \in \text{Paths}(s)$

with  $\pi \models \Diamond T$  we have  $\pi \models \Box \Diamond G$ .

By the law of large numbers theorem

$$\Pr_s \{ \pi \in \text{Paths}(s) \mid \pi \models \Box \Diamond G \} = 1 \quad \square.$$

# Almost sure repeated reachability

## Almost sure repeated reachability is PCTL-definable

For finite DTMC  $\mathcal{D}$ , state  $s \in S$  and  $G \subseteq S$ :

$$s \models \mathbb{P}_{=1}(\Box \mathbb{P}_{=1}(\Diamond G)) \quad \text{iff} \quad Pr_s\{\pi \in Paths(s) \mid \pi \models \Box \Diamond G\} = 1.$$

We abbreviate  $\mathbb{P}_{=1}(\Box \mathbb{P}_{=1}(\Diamond G))$  by  $\mathbb{P}_{=1}(\Box \Diamond G)$ .

### Proof:

On the blackboard.

### Remark:

For CTL, universal repeated reachability properties can be formalized by the combination of the modalities  $\forall \Box$  and  $\forall \Diamond$ :

$$s \models \forall \Box \forall \Diamond G \quad \text{iff} \quad \pi \models \Box \Diamond G \text{ for all } \pi \in Paths(s).$$

$\forall \Diamond \forall \Box G$

$\neq$

$\Box \Diamond G$




# Repeated reachability probabilities

## Repeated reachability probabilities are PCTL-definable

For finite DTMC  $\mathcal{D}$ , state  $s \in S$ ,  $G \subseteq S$  and interval  $J \subseteq [0, 1]$  we have:

$$\underbrace{s \models \underbrace{\mathbb{P}_J(\diamond \mathbb{P}_{=1}(\square \mathbb{P}_{=1}(\diamond G)))}_{=\mathbb{P}_J(\square \diamond G)}}_{\text{PCTL}} \quad \text{if and only if} \quad \underbrace{Pr(s \models \square \diamond G)}_{=1} \in J.$$

PCTL



$$Pr(s \models \square \diamond G) \in J$$

# Repeated reachability probabilities

## Repeated reachability probabilities are PCTL-definable

For finite DTMC  $\mathcal{D}$ , state  $s \in S$ ,  $G \subseteq S$  and interval  $J \subseteq [0, 1]$  we have:

$$s \models \underbrace{\mathbb{P}_J(\Diamond \mathbb{P}_{=1}(\Box \mathbb{P}_{=1}(\Diamond G)))}_{=\mathbb{P}_J(\Box \Diamond G)} \quad \text{if and only if} \quad \Pr(s \models \Box \Diamond G) \in J.$$

### Proof:

By the long run theorem (cf. Lecture 4), almost surely a BSCC  $T$  will be reached and each of its states will be visited infinitely often. Thus, the probabilities for  $\Box \Diamond G$  agree with the probability to reach a BSCC  $T$  that contains a state in  $G$ .

### Remark:

By the above theorem,  $\mathbb{P}_{>0}(\Box \Diamond G)$  is *qualitative* PCTL definable. Note that  $\exists \Box \Diamond G$  is not CTL-definable (but definable in a combination of CTL and LTL, called CTL\*).

# Almost sure persistence

$$P_{=1}(\Diamond\Box G) \approx \forall\Diamond\Box G \neq \Diamond\Box G$$

# Almost sure persistence

## Almost sure persistence is PCTL-definable

For finite DTMC  $\mathcal{D}$ , state  $s \in S$  and  $G \subseteq S$ :

$$s \models \mathbb{P}_{=1}(\Diamond \mathbb{P}_{=1}(\Box G)) \quad \text{iff} \quad \underbrace{Pr_s\{\pi \in Paths(s) \mid \pi \models \Diamond \Box G\}}_{\text{almost surely}} = 1.$$

$$s \models \Diamond \Box G$$

# Almost sure persistence

## Almost sure persistence is PCTL-definable

For finite DTMC  $\mathcal{D}$ , state  $s \in S$  and  $G \subseteq S$ :

$$s \models \mathbb{P}_{=1}(\Diamond \mathbb{P}_{=1}(\Box G)) \quad \text{iff} \quad Pr_s\{\pi \in Paths(s) \mid \pi \models \Diamond \Box G\} = 1.$$

We abbreviate  $\mathbb{P}_{=1}(\Diamond \mathbb{P}_{=1}(\Box G))$  by  $\mathbb{P}_{=1}(\Diamond \Box G)$ .

### Proof:

Left as an exercise.

### Remark:

Note that  $\forall \Diamond \Box G$  is not CTL-definable.  $\Diamond \Box G$  is a well-known example formula in LTL that cannot be expressed in CTL. But by the above theorem it can be expressed in PCTL.

# Persistence probabilities

## Persistence probabilities are PCTL-definable

For finite DTMC  $\mathcal{D}$ , state  $s \in S$ ,  $G \subseteq S$  and interval  $J \subseteq [0, 1]$  we have:

$$s \models \underbrace{\mathbb{P}_J(\Diamond \mathbb{P}_{=1}(\Box G))}_{=\mathbb{P}_J(\Diamond \Box G)} \quad \text{if and only if} \quad \Pr(s \models \Diamond \Box G) \in J.$$

## Proof:

Left as an exercise. Hint: use the long run theorem (cf. Lecture 4).

# Overview

- 1 Introduction
- 2 Qualitative PCTL
- 3 Computation Tree Logic
- 4 CTL versus qualitative PCTL
- 5 Fair CTL versus qualitative PCTL
- 6 Repeated reachability and persistence
- 7 Summary**

# Summary

- ▶ Qualitative PCTL only allow the probability bounds  $> 0$  and  $= 1$ .
- ▶ There is no CTL formula that is equivalent to  $\mathbb{P}_{=1}(\Diamond a)$ .
- ▶ There is no PCTL formula that is equivalent to  $\forall \Box a$ .
- ▶ These results do not apply to finite DTMCs.
- ▶  $\mathbb{P}_{=1}(\Diamond a)$  and  $\forall \Diamond a$  are equivalent under strong fairness.
- ▶ Repeated reachability probabilities are PCTL definable.

## Take-home messages

Qualitative PCTL and CTL have incomparable expressiveness. Qualitative and fair CTL are equally expressive. Repeated reachability and persistence probabilities are PCTL definable. Their qualitative counterparts are not all expressible in CTL.