

Probabilistic Programming

Lecture #5: Domain Theory

Joost-Pieter Katoen



RWTH Lecture Series on Probabilistic Programming 2018

Overview

- 1 Motivation
- 2 Complete lattices
- 3 Monotonic and continuous functions
- 4 Fixpoint theorems

Overview

- 1 Motivation
- 2 Complete lattices
- 3 Monotonic and continuous functions
- 4 Fixpoint theorems

Aims and sufficient conditions

- ▶ In denotational program semantics, the semantics of a loop is defined as some **fixed point** of a mathematical function
 - ▶ We will consider this for pGCL
- ▶ **Goals:**
 - ▶ Prove **existence** of such fixed points
 - ▶ Show how they can be “computed” (more exactly: **approximated**)
- ▶ **Sufficient conditions:**
 - ▶ on function domains: complete lattices
 - ▶ on functions: monotonicity and Scott continuity

Overview

- 1 Motivation
- 2 Complete lattices
- 3 Monotonic and continuous functions
- 4 Fixpoint theorems

Upper and lower bounds

Upper bound

An element $d \in D$ is called an **upper bound** of $S \subseteq D$, denoted $S \sqsubseteq d$, if $s \sqsubseteq d$ for every $s \in S$.

An upper bound d of $S \subseteq D$ is called **least upper bound (LUB)** or **supremum** of S , denoted $d = \sqcup S$, if $d \sqsubseteq d'$ for every upper bound d' of S .

Lower bound

An element $d \in D$ is called a **lower bound** of $S \subseteq D$, denoted $d \sqsupseteq S$, if $d \sqsupseteq s$ for every $s \in S$.

A lower bound d of $S \subseteq D$ is called **greatest lower bound (GLB)** or **infimum** of S , denoted $d = \sqcap S$, if $d' \sqsupseteq d$ for every lower bound d' of S .

Partial orders

Partial order

A **partial order (PO)** (D, \sqsubseteq) consists of a set D , called **domain**, and of a relation $\sqsubseteq \subseteq D \times D$ such that, for every $d_1, d_2, d_3 \in D$:

- ▶ $d_1 \sqsubseteq d_1$ (reflexivity)
- ▶ $d_1 \sqsubseteq d_2$ and $d_2 \sqsubseteq d_3 \implies d_1 \sqsubseteq d_3$ (transitivity)
- ▶ $d_1 \sqsubseteq d_2$ and $d_2 \sqsubseteq d_1 \implies d_1 = d_2$ (antisymmetry)

A PO is called **total** if, in addition, always $d_1 \sqsubseteq d_2$ or $d_2 \sqsubseteq d_1$.

Examples

1. (\mathbb{N}, \leq) is a total partial order
2. $(2^{\mathbb{N}}, \subseteq)$ is a (non-total) partial order
3. $(\mathbb{N}, <)$ is not a partial order (since not reflexive)

Chains

Chains

$S \subseteq D$ is called a **chain** in D if, for every $s_1, s_2 \in S$,

$$s_1 \sqsubseteq s_2 \text{ or } s_2 \sqsubseteq s_1.$$

That is, S is a totally ordered subset of D .

A chain $S = s_1 \sqsubseteq s_2 \sqsubseteq s_3 \sqsubseteq \dots$ is a **ascending**.


A chain $S = s_1 \sqsupseteq s_2 \sqsupseteq s_3 \sqsupseteq \dots$ is a **descending**.

Examples: chains

- Every subset $S \subseteq \mathbb{N}$ is a chain in (\mathbb{N}, \leq) .
It has a LUB (its greatest element) iff it is finite.
- $\{\emptyset, \{0\}, \{0, 1\}, \dots\}$ is a chain in $(2^{\mathbb{N}}, \subseteq)$ with LUB \mathbb{N} .

Examples: lattices

- $(2^{\mathbb{N}}, \subseteq)$ is a complete lattice with $\bigsqcup S = \bigcup_{M \in S} M$ for every subset $S \subseteq 2^{\mathbb{N}}$.
- (\mathbb{N}, \leq) is not a complete lattice, as, e.g., the chain \mathbb{N} has no upper bound.
- Which of the following structures are complete lattices?



Complete lattices

Complete lattice

A PO (D, \sqsubseteq) is called a **complete lattice**, if every subset $S \subseteq D$ has a supremum in D , i.e., $\bigsqcup S \in D$.

Every complete lattice (D, \sqsubseteq) has a **least** element \perp and, dually, a **greatest** element \top which satisfy:

$$\forall d \in D. \perp \sqsubseteq d \sqsubseteq \top.$$

For complete lattice (D, \sqsubseteq) : every subset $S \subseteq D$ has an infimum in D , i.e., $\bigsqcap S \in D$.

Every ascending or descending chain has a least upper bound and greatest lower bound.

Overview

- Motivation
- Complete lattices
- Monotonic and continuous functions
- Fixpoint theorems

Monotonicity

Monotonicity

Let (D, \sqsubseteq) and (D', \sqsubseteq') be partial orders, and let $F : D \rightarrow D'$. F is called **monotonic** (w.r.t. (D, \sqsubseteq) and (D', \sqsubseteq')) if, for every $d_1, d_2 \in D$,

$$d_1 \sqsubseteq d_2 \text{ implies } F(d_1) \sqsubseteq' F(d_2).$$

Interpretation: monotonic functions **preserve information**

1. Let $T := \{S \subseteq \mathbb{N} \mid S \text{ finite}\}$. Then $F_1 : T \rightarrow \mathbb{N} : S \mapsto \sum_{n \in S} n$ is monotonic w.r.t. $(2^{\mathbb{N}}, \sqsubseteq)$ and (\mathbb{N}, \leq) .
2. $F_2 : 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}} : S \mapsto \mathbb{N} \setminus S$ is not monotonic w.r.t. $(2^{\mathbb{N}}, \sqsubseteq)$ (since, e.g., $\emptyset \subseteq \mathbb{N}$ but $F_2(\emptyset) = \mathbb{N} \not\subseteq F_2(\mathbb{N}) = \emptyset$).

Scott continuity

A function F is continuous if applying F and taking LUBs is commutable:

Scott continuity

Let (D, \sqsubseteq) and (D', \sqsubseteq') be complete lattices and $F : D \rightarrow D'$ monotonic. Then F is called **continuous** if, for every non-empty chain $S \subseteq D$,

$$F\left(\bigsqcup S\right) = \bigsqcup F(S).$$

Every continuous function is monotonic.

Proof.

1. Let $d \sqsubseteq e$. Then $\{d, e\}$ is a chain with $\bigsqcup\{d, e\} = e$.
2. Let F be continuous. Then $F(d) \sqsubseteq \bigsqcup\{F(d), F(e)\}$.
3. By continuity of F , $\{F(d), F(e)\} = F(\bigsqcup\{d, e\})$, which equals $F(d)$.

Monotonicity on chains

The following lemma states how chains behave under monotonic functions.

Let (D, \sqsubseteq) and (D', \sqsubseteq') be complete lattices, $F : D \rightarrow D'$ monotonic, and $S \subseteq D$ a chain in D . Then:

1. $F(S) := \{F(d) \mid d \in S\}$ is a chain in D' .
2. $\bigsqcup F(S) \sqsubseteq' F(\bigsqcup S)$.

Proof.

Left as a homework exercise. □

Overview

- 1 Motivation
- 2 Complete lattices
- 3 Monotonic and continuous functions
- 4 Fixpoint theorems

Fixed points


Fixed point

Let $F : D \rightarrow D$ be a function. Element $d \in D$ is called a **fixed point** of F if and only if $F(d) = d$.

Examples

1. Function $F : \mathbb{R} \rightarrow \mathbb{R}$ with $F(x) = x^2 + 3x + 4$ has a fixed point at 2.
2. For function $F : \mathbb{R} \rightarrow \mathbb{R}$ with $F(x) = x$, all $x \in \mathbb{R}$ are fixed points.
3. Function $F : \mathbb{R} \rightarrow \mathbb{R}$ with $F(x) = x+4$ has no fixed points.
4. Function $F : \mathbb{R} \rightarrow \mathbb{R}$ with $F(x) = \frac{x}{2} + \frac{1}{x}$ has a fixed point at $\sqrt{2}$.
5. Function $F : \mathbb{R} \rightarrow \mathbb{R}$ with $F(x) = \cos(x)$ has a fixed point, but this is hard to determine.

Iterative scheme to determine a fixed point



How to find fixed points?

Naive scheme: start with an initial value x_0 , and then iterate: $x_{n+1} = f(x_n)$.
 (Wrong) idea: as n grows larger, x_n converges to some fixed point of F .

1. Take $F(x) = \frac{x}{2} + \frac{1}{x}$ and $x_0 = 1$. Iterations yields: $\frac{3}{2}, \frac{17}{12}, \frac{17}{24} + \frac{12}{17}$ which indeed approximates $\sqrt{2}$.
2. Take $F(x) = \frac{5}{2}x - \frac{3}{2}x^2$. Iteration converges to 1.
3. But, take $F(x) = \frac{13}{4}x - \frac{3}{2}x^2$. Iteration oscillates between two points, regardless of the initial value.

When does such an iterative scheme (i.e., approximate) a fixed point, and if so, which fixed point?

We consider this for complete lattices and continuous functions.

Kleene's fixpoint theorem

Kleene's fixpoint theorem

Let (D, \sqsubseteq) be a complete lattice and $F : D \rightarrow D$ continuous. Then F has a least fixed point $\text{lfp } F$ and greatest fixed point $\text{gfp } F$ respectively, given by:

$$\text{lfp } F := \sup_{n \in \mathbb{N}} F^n(\perp) \quad \text{and} \quad \text{gfp } F := \inf_{n \in \mathbb{N}} F^n(\top)$$

where $F^0(d) = d$ and $F^{n+1}(d) = F(F^n(d))$.

Proof.

on the board



Examples

- ▶ **Domain:** complete lattice $(2^{\mathbb{N}}, \subseteq)$ with $\bigsqcup S = \bigcup_{N \in S} N$
- ▶ **Function:** $F : 2^{\mathbb{N}} \rightarrow 2^{\mathbb{N}} : N \mapsto N \cup A$ for some fixed $A \subseteq \mathbb{N}$
 - ▶ F monotonic: $M \subseteq N \implies F(M) = M \cup A \subseteq N \cup A = F(N)$
 - ▶ F continuous: $F(\bigsqcup S) = F(\bigcup_{N \in S} N) = (\bigcup_{N \in S} N) \cup A = \bigcup_{N \in S} (N \cup A) = \bigcup_{N \in S} F(N) = \bigsqcup F(S)$.
- ▶ **Fixpoint iteration:** $N_n := F^n(\bigsqcup \emptyset)$ where $\bigsqcup \emptyset = \emptyset$
 - ▶ $N_0 = \bigsqcup \emptyset = \emptyset$
 - ▶ $N_1 = F(N_0) = \emptyset \cup A = A$
 - ▶ $N_2 = F(N_1) = A \cup A = A = N_n$ for every $n \geq 1$ $\implies \text{gfp } F = A$
- ▶ **Alternatively:** $F(N) = N \cap A$
 $\implies \text{gfp } F = \emptyset$

Knaster-Tarski theorem (2)

Knaster-Tarski theorem

For any complete lattice (D, \subseteq) the following holds:

1. The least fixed and the prefixed points of F exist, and are identical
2. The greatest fixed and the postfix points of F exist, and are identical
3. The fixed points of F form a complete lattice.

Proof.

On the black board. □

Knaster-Tarski theorem (1)

Alfred Tarski (1901–1983)



Bronislaw Knaster (1893–1990)



Prefix point and postfix point

For monotonic function $F : D \rightarrow D$ on (D, \subseteq) and $d \in D$:

1. d is a prefixed point of F if $F(d) \subseteq d$
2. d is a postfix point of F if $d \subseteq F(d)$.

d is a prefixed and a postfix point of F iff d is a fixed point.

Pictorial depiction of the Knaster-Tarski theorem

